

New Jersey Division of HIV/AIDS Services  
Persons Living with HIV/AIDS as of June 30, 2004  
Willingboro Residents

1. Current HIV/AIDS Category	Adult/Adolescent *		Pediatric *		Total	
	Cases	%	Cases	%	Cases	%
-----	-----	---	-----	---	-----	---
AIDS-Any OI	#	#	#	#	12	12
AIDS-CD4 Only	#	#	#	#	32	33
HIV Only	54	56	0	0	54	55
-----	-----	---	-----	---	-----	---
Total	97	100	#	#	98	100

Current			3. Race/Eth.	Adult/Adl*		Pediatric*		Total	
Age *	Cases	%		Cases	%	Cases	%	Cases	%
-----	-----	---	-----	-----	---	-----	---	-----	---
<5	0	0	White, Not Hispanic	10	10	0	0	10	10
5-12	#	#	Black, Not Hispanic	#	#	#	#	81	83
13-19	#	#	Hispanic	5	5	0	0	5	5
20-29	10	10	Other/Unknown	#	#	#	#	#	#
30-39	25	26	-----	-----	---	-----	---	-----	---
40-49	38	39	Total	97	100	#	#	98	100
50+	24	24							
Unk.	0	0							
-----	-----	---							
Total	98	100							

4. Patient Group	Adult/Adolescent Transmission Modes **					
	Males	%	Females	%	Total	%
-----	-----	---	-----	---	-----	---
Men sex with men (MSM)	29	44	0	0	29	30
Injection drug user (IDU)	#	#	#	#	#	#
MSM/IDU	#	#	0	0	#	#
Heterosexual contact	#	#	#	#	17	18
Risk not reported/other***	17	26	15	48	32	33
-----	-----	---	-----	---	-----	---
Total	66	100	31	100	97	100

	Pediatric Transmission Modes					
	Males	%	Females	%	Total	%
-----	-----	---	-----	---	-----	---
Parent at risk/has AIDS/HIV	#	#	#	#	#	#
Risk not reported/other***	0	0	0	.	0	0
-----	-----	---	-----	---	-----	---
Total	#	#	#	#	#	#

NOTE: Includes only those persons known to be infected with HIV.

# Indicates that number is not shown due to small cell size,  
in accordance with NJDHSS security and confidentiality policies.

\* Classification based on current age at the end of June 2004.

\*\* 0 patients are currently of adult/adolescent age (13+ years old), but have evidence of having been HIV-infected as children. They are counted as adult/adolescent reports in tables 1,2, and 3, and as pediatric reports in table 4.

\*\*\* Included in this category are persons with modes of transmission of coagulation disorder and transfusion with blood/blood products.

New Jersey Division of HIV/AIDS Services  
AIDS Cases Reported as of June 30, 2004  
Willingboro Residents

1. Disease Category	Adult/Adolescent*		Deaths (%)	Pediatric*		Deaths (%)	Total	
	Cases	(%)		Cases	(%)		Cases	(%)
PCP	41	( 34)	35 ( 88)	0	( 0)	0 ( .)	41	( 34)
Other OI w/o PCP	#	( #)	# ( #)	#	( #)	# ( #)	#	( #)
KS Alone	#	( #)	# ( #)	#	( #)	# ( #)	#	( #)
No Disease Listed	42	( 35)	10 ( 38)	0	( 0)	0 ( .)	42	( 35)
Total	119	(100)	76 ( 76)	#	( #)	# ( #)	121	(100)

2. Age *	Cases		3. Race/Eth.	Adult/Adl*		Pediatric*		Total	
	Cases	%		Cases	%	Cases	%	Cases	%
Under 5	#	#	White, Not Hispanic	#	#	#	#	15	12
5-12	#	#	Black, Not Hispanic	#	#	#	#	98	81
13-19	0	0	Hispanic	8	7	0	0	8	7
20-29	25	21	Other/Unknown	0	0	0	0	0	0
30-39	45	37	-----	-----	---	-----	---	-----	---
40-49	35	29	Total	119	100	#	#	121	100
Over 49	14	12							
Unknown	0	0							
Total	121	100							

4. Patient Group	Adult/Adolescent Transmission Modes **					
	Males	%	Females	%	Total	%
Men sex with men (MSM)	44	46	0	0	44	37
Injection drug user (IDU)	25	26	8	33	33	28
MSM/IDU	7	7	0	0	7	6
Heterosexual contact	6	6	8	33	14	12
Risk not reported/other***	13	14	8	33	21	18
Total	95	100	24	100	119	100

	Pediatric Transmission Modes					
	Males	%	Females	%	Total	%
Parent at risk/has AIDS/HIV	#	#	#	#	#	#
Risk not reported/other***	0	0	0	0	0	0
Total	#	#	#	#	#	#

# indicates that number is not shown due to small cell size,  
in accordance with NJDHSS security and confidentiality policies.

\* Classification at time of AIDS diagnosis.

\*\* 0 patients were diagnosed with AIDS as adults, but have evidence either of having been HIV-infected or of having acquired AIDS as children. They are counted as adult/adolescent reports in tables 1,2, and 3; and as pediatric reports in table 4.

\*\*\* Included in this category are persons with modes of transmission of coagulation disorder and transfusion with blood/blood products.

New Jersey Division of HIV/AIDS Services  
AIDS Cases Reported as of June 30, 2004  
Willingboro Residents

5. AIDS Cases by Year of Diagnosis.

Year of Diagnosis -----	Number of Cases -----	Number of Deaths -----	Case-Fatality Rate -----
Before 1985	#	#	#
1985	#	#	#
1986	#	#	#
1987	#	#	#
1988	10	9	90%
1989	11	11	100%
1990	9	9	100%
1991	10	10	100%
1992	9	9	100%
1993	7	6	86%
1994	14	8	57%
1995	7	#	#
1996	5	#	#
1997	#	#	#
1998	5	#	#
1999	7	#	#
2000	#	#	#
2001	#	#	#
2002	8	0	0%
2003	5	0	0%
2004	#	#	#
-----	-----	-----	-----
Total	121	77	64%

New Jersey Division of HIV/AIDS Services  
HIV INFECTION-not-AIDS Reports with Identifiers Reported as of June 30, 2004  
Willingboro Residents

1. Age *	Cases		2. Race/Ethnicity	Adult/Adl*		Pediatric*		Total	
	-----	---		Cases	%	Cases	%	Cases	%
-----	-----	---	-----	-----	---	-----	---	-----	---
Under 5	0	0	White, Not Hispanic	#	#	#	#	#	#
5-12	0	0	Black, Not Hispanic	49	82	0	.	49	83
13-19	#	#	Hispanic	0	0	0	.	0	0
20-29	#	#	Other/Unknown	#	#	0	.	#	#
30-39	17	28	-----	-----	---	-----	---	-----	---
40-49	15	25	Total	60	100	#	#	60	100
Over 49	8	13							
Unknown	0	0							
-----	-----	---							
Total	60	100							

3. Patient Group	Adult/Adolescent Transmission Modes **					
	Males	%	Females	%	Total	%
-----	-----	---	-----	---	-----	---
Men sex with men (MSM)	18	49	0	0	18	30
Injection drug user (IDU)	6	16	5	22	11	18
MSM/IDU	#	#	0	0	#	#
Heterosexual contact	#	#	#	#	#	#
Risk not reported/other***	9	24	11	48	20	33
-----	-----	---	-----	---	-----	---
Total	37	100	23	100	60	100

	Pediatric Transmission Modes				Total	%
	Males	%	Females	%		
-----	-----	---	-----	---	-----	---
Parent at risk/has AIDS/HIV	0	.	0	.	0	.
Risk not reported/other***	0	.	0	.	0	.
-----	-----	---	-----	---	-----	---
Total	0	.	0	.	0	.

\* Classification at time of first HIV+ test or at time of report if not yet confirmed HIV+.

# indicates that number is not shown due to small cell size, in accordance with NJDHSS security and confidentiality policies.

\*\* 0 patients were diagnosed with HIV as adults, but have evidence of having been HIV-infected as children. They are counted as adult/adolescent reports in tables 1 and 2; and as pediatric reports in table 3.

\*\*\* Included in this category are persons with modes of transmission of coagulation disorder and transfusion with blood/blood products.

NOTE: These tables include those reports of HIV Positive infection that have not yet been updated to AIDS status. Not included in these tables are cases initially reported as HIV-not-AIDS that have subsequently been updated to AIDS. Also not included in these tables are HIV Perinatal Exposure Reports where the serostatus of the child is not confirmed HIV Positive; i.e., the HIV serostatus of the child is undetermined or has been found to be negative (seroreverted).

New Jersey Division of HIV/AIDS Services  
HIV/AIDS Cases Reported as of June 30, 2004  
Willingboro Residents

1. Current HIV/AIDS Category	Adult/Adolescent *		Pediatric *		Total	
	Cases	%	Cases	%	Cases	%
-----	----	---	----	---	----	---
AIDS-Any OI	#	#	#	#	79	44
AIDS-CD4 Only	#	#	#	#	42	23
HIV Only	60	34	0	0	60	33
-----	----	---	----	---	----	---
Total	179	100	#	#	181	100

2. Age *	Cases	%	3. Race/Eth.	Adult/Adl*		Pediatric*		Total	
-----	-----	---	-----	Cases	%	Cases	%	Cases	%
Under 5	#	#	White, Not Hispanic	24	13	0	0	24	13
5-12	0	0	Black, Not Hispanic	#	#	#	#	147	81
13-19	5	3	Hispanic	8	4	0	0	8	4
20-29	44	24	Other/Unknown	#	#	#	#	#	#
30-39	63	35	-----	-----	---	-----	---	-----	---
40-49	45	25	Total	179	100	#	#	181	100
Over 49	22	12							
Unknown	#	#							
-----	-----	---							
Total	181	100							

4. Patient Group	Adult/Adolescent Transmission Modes **					
	Males	%	Females	%	Total	%
-----	-----	---	-----	---	-----	---
Men sex with men (MSM)	62	47	0	0	62	35
Injection drug user (IDU)	31	23	13	28	44	25
MSM/IDU	8	6	0	0	8	4
Heterosexual contact	9	7	15	32	24	13
Risk not reported/other***	22	17	19	40	41	23
-----	-----	---	-----	---	-----	---
Total	132	100	47	100	179	100

	Pediatric Transmission Modes					
	Males	%	Females	%	Total	%
-----	-----	---	-----	---	-----	---
Parent at risk/has AIDS/HIV	#	#	#	#	#	#
Risk not reported/other***	0	0	0	0	0	0
-----	-----	---	-----	---	-----	---
Total	#	#	#	#	#	#

NOTE: Includes only those persons known to be infected with HIV.

# Indicates that number is not shown due to small cell size,  
in accordance with NJDHSS security and confidentiality policies.

\* Classification at time of first HIV+/AIDS diagnosis.

\*\* 0 patients were diagnosed with HIV/AIDS as adults, but have evidence of having  
been HIV-infected as children. They are counted as adult/adolescent  
reports in tables 1,2 and 3; and as pediatric reports in table 4.

\*\*\* Included in this category are persons with modes of transmission of  
coagulation disorder and transfusion with blood/blood products.

New Jersey Division of HIV/AIDS Services  
HIV/AIDS Cases Reported as of June 30, 2004  
Willingboro Residents

5. HIV/AIDS Cases by Year of First HIV/AIDS Diagnosis.

Year of First HIV/AIDS Diagnosis	Number of Cases	Number Known Dead	Case-Fatality Rate
-----	-----	-----	-----
Before 1985	0	0	#
1985	#	#	#
1986	#	#	#
1987	6	5	83%
1988	12	11	92%
1989	15	13	87%
1990	10	8	80%
1991	9	8	89%
1992	17	10	59%
1993	13	#	#
1994	11	5	45%
1995	15	5	33%
1996	6	#	#
1997	10	#	#
1998	12	#	#
1999	8	#	#
2000	5	0	0%
2001	5	0	0%
2002	9	0	0%
2003	9	0	0%
2004	#	#	#
-----	-----	-----	-----
Total	181	83	46%